



SLAP REPAIR PROTOCOL

Name: _____

Diagnosis: _____

Date of Surgery: _____

Frequency: 1 2 3 4 times / week Duration: 1 2 3 4 5 6 Weeks

Week 0---1: Patient to do Home Exercises (pendulums, elbow ROM, wrist ROM, grip strengthening)

____ Weeks 1--4:

- No IR up the back; No ER behind the head
- ROM goals: 90° FF/20° ER at side
- No resisted FF or biceps until 6 weeks post---op as to not stress the biceps root
- Sling for 4 weeks
- Heat before/ice after PT sessions

____ Weeks 4--8:

- D/C sling
- Increase AROM 140° FF/ 40° ER at side/ 60° ABD/ IR behind back to waist
- Strengthening (isometrics/light bands) within AROM limitations
- Also start strengthening scapular stabilizers (traps/rhomboids/lev. scap/etc)
- Physical modalities per PT discretion

____ Weeks 8--12:

- If ROM lacking, increase to full with gentle passive stretching at end ranges
- Advance strengthening as tolerated: isometrics → bands → light weights (1---5 lbs); 8---12 reps/2--3 set per rotator cuff, deltoid, and scapular stabilizers

____ Months 3---12:

- Only do strengthening 3x/week to avoid rotator cuff tendonitis
- Begin UE ergometer
- Begin eccentrically resisted motions, plyometrics (ex weighted ball toss), proprioception (ex body blade), and closed chain exercises at 12 weeks.
- Begin sports related rehab at 3 months, including advanced conditioning
- Return to throwing at 4 1/2 months
- Throw from pitcher's mound at 6 months

Modalities/Other:

Signature _____

Date: _____